

# CONFORMAL EQUIVALENCE BETWEEN CERTAIN GEOMETRIES IN DIMENSION 6 AND 7

RICHARD CLEYTON AND STEFAN IVANOV

**ABSTRACT.** For  $G_2$ -manifolds the Fernández-Gray class  $\mathcal{X}_1 + \mathcal{X}_4$  is shown to consist of the union of the class  $\mathcal{X}_4$  of  $G_2$ -manifolds locally conformal to parallel  $G_2$ -structures and that of conformal transformations of nearly parallel or weak holonomy  $G_2$ -manifolds of type  $\mathcal{X}_1$ . The analogous conclusion is obtained for Gray-Hervella class  $\mathcal{W}_1 + \mathcal{W}_4$  of real 6-dimensional almost Hermitian manifolds: this sort of geometry consists of locally conformally Kähler manifolds of class  $\mathcal{W}_4$  and conformal transformations of nearly Kähler manifolds in class  $\mathcal{W}_1$ . A corollary of this is that a compact  $SU(3)$ -space in class  $\mathcal{W}_1 + \mathcal{W}_4$  or  $G_2$ -space of the kind  $\mathcal{X}_1 + \mathcal{X}_4$  has constant scalar curvature if only if it is either a standard sphere or a nearly parallel  $G_2$  or nearly Kähler manifold, respectively. The properties of the Riemannian curvature of the spaces under consideration are also explored.

## 1. INTRODUCTION

Reductions of the bundle of orthonormal frames over a Riemannian manifold to a principal  $G$ -bundle may be classified by the  $G$ -invariant components of the intrinsic torsion.

This idea was originally due to Gray and collaborators [10, 17] for the special instances of  $G_2$ -manifolds and almost Hermitian manifolds. It has been further refined and explored by, for instance, Bryant [6], Farinola, Falcitelli & Salamon [9], Martín Cabrera [24, 23], Martín Cabrera, Monar & Swann [25], Chiossi & Salamon [8].

For  $G_2$ - and almost Hermitian structures alike, the intrinsic torsion has 4 irreducible components. There are thus potentially 16 torsion classes for these two kinds of geometries.

In [25], Martín Cabrera, Monar & Swann showed that apart from one instance,  $\mathcal{X}_1 + \mathcal{X}_2$  in our notation, every single class of  $G_2$ -structures may be realized on a compact homogeneous space. For the one exception an easy calculation shows that any  $G_2$  structure with torsion  $X_1 + X_2$  must have either  $X_1 = 0$  or  $X_2 = 0$ .

In section 3 we show that something similar holds for the class  $\mathcal{X}_1 + \mathcal{X}_4$ . Namely that the latter essentially is generated by the classes  $\mathcal{X}_1$  and  $\mathcal{X}_4$ . It is well known that  $G_2$ -structures in this class are locally conformally equivalent to nearly parallel ones. We will show that this equivalence is only really local when the  $G_2$ -structure lies in the subclass  $\mathcal{X}_4$  of locally conformally parallel structures. The structure of compact locally conformally parallel  $G_2$ -manifolds has been recently described in [20, 29]. In contrast to this, we will show that if the  $X_1$  component is non-zero at some point, it is non-zero everywhere. This is the key point in proving that a global conformal change exists.

Differently from the  $G_2$  case, it was only recently pointed out by Butruille [7] that a 6-dimensional almost Hermitian manifold in the Gray-Hervella class  $\mathcal{W}_1 + \mathcal{W}_4$  is locally conformal to a nearly Kähler manifold. In section 5, we present a different and simpler proof of this fact for completeness. Based on this the analogous statements to those given for  $G_2$ -structures are shown to hold for 6-dimensional almost Hermitian geometry, too. In particular, any almost Hermitian 6-manifold in the class  $(\mathcal{W}_1 + \mathcal{W}_4) \setminus \mathcal{W}_4$  has trivial canonical bundle. The geometries  $\mathcal{W}_4$  for  $SU(3)$ - and  $\mathcal{X}_4$  for  $G_2$ -manifolds are both special instances of  $G$ -structures with vectorial torsion. This notion was studied in [1]. The almost Hermitian manifolds and  $G_2$ -structures studied in this paper all fit in the wider framework of  $G$ -structures with three-form torsion, see [12, 2].

The aim of this note is to establish

**Theorem 1.** *Let  $(M, g, \phi)$  be a compact 7-dimensional manifold locally conformally equivalent to a nearly parallel  $G_2$ -manifold. Then  $(M, g, \phi)$  has constant scalar curvature if and only if  $(M, g)$  is either nearly parallel or conformally equivalent to the standard 7-sphere.*

**Theorem 2.** *Let  $(M, g, J)$  be a compact 6-dimensional manifold locally conformally equivalent to a nearly Kähler manifold. Then  $(M, g, J)$  has constant scalar curvature if and only if  $(M, g)$  is either nearly parallel or conformally equivalent to the 6-sphere with standard metric.*

In the last section we characterize complete Einstein  $G_2$  and  $SU(3)$  manifolds in the strict class  $\mathcal{X}_1 + \mathcal{X}_4$  and  $\mathcal{W}_1 + \mathcal{W}_4$ , respectively. The phrases ‘strict class’ is used here to indicate that the  $G$ -structure is not in any sub-class of the one given. So a  $G_2$ -structure strictly in class  $\mathcal{X}_1$  must, in particular, have non-trivial intrinsic torsion.

The results obtained in this paper are direct consequences of the following. The  $G$ -structures under consideration are described by the existence of certain *fundamental differential forms*  $\omega_1, \dots, \omega_p$ , whose exterior derivatives determine the corresponding intrinsic torsion in full. First order identities on the  $G$ -invariant components of the intrinsic torsion descend from the closure of  $d\omega_i$ . These equations in general have non-trivial consequences as is seen by the examples considered here.

The relations coming from the second derivatives of the forms may also be seen as consequences of the first Bianchi identity, see for instance [26, 6].

*Acknowledgements.* The authors wish to thank Andrew Swann, Ilka Agricola and Thomas Friedrich for helpful conversations and Simon Chiossi for many helpful comments. The first named author was supported by the Junior Research Group “Special Geometries in Mathematical Physics”, of the Volkswagen foundation and the SFB 647 “Space–Time–Matter”, of the DFG. The authors wish to thank UC Riverside for hospitality and Yat Sun Poon for support during the initial stage of this project.

## 2. A LEMMA

The key to obtaining the results is the observation

**Lemma 3.** *Let  $M$  be a connected manifold equipped with a differentiable function  $\phi \not\equiv 0$  and a one-form  $\alpha$  such that*

$$(2.1) \quad \begin{aligned} d\alpha &= 0, \\ d\phi + \phi\alpha &= 0. \end{aligned}$$

*Then  $\phi$  is nowhere zero and  $\alpha = -d \log |\phi|$ .*

*Proof.* Let  $\phi$  and  $\alpha$  be a function and one-form as in equation (2.1). By Poincaré’s Lemma we may choose a covering  $U_i$  of  $M$  and functions  $f_i: U_i \rightarrow \mathbb{R}$  such that  $\alpha|_{U_i} = df_i$ . Then equation (2.1) implies that the product  $\phi \exp(f_i)$  is constant over each  $U_i$ . Therefore, if  $\phi(p) \neq 0$  at some point  $p$  in, say  $U_0$ , then  $\phi \neq 0$  over all  $U_0$  and therefore also on each  $U_j$  that overlaps  $U_0$ . The conclusion now follows from connectedness.  $\square$

## 3. THE $G_2$ CASE

A  $G_2$ -manifold is a 7-dimensional manifold  $M$  equipped with a special, so-called fundamental three-form  $\phi$ , required to satisfy the following non-degeneracy condition

$$(3.1) \quad i_X \phi \wedge i_Y \phi \wedge \phi = 6g(X, Y) \text{ vol}(g),$$

for some Riemannian metric  $g$  and orientation on  $M$ . The notation  $i_X \phi$  means interior product of the vector field  $X$  with the three-form  $\phi$ . It is well known that the covariant derivative of the fundamental three-form is determined by the exterior derivatives of  $\phi$  and its Hodge dual  $*\phi$ . Using the representation theory of  $G_2$  on the exterior algebra one may write these differentials as

$$\begin{aligned} d\phi &= \tau_0 * \phi + 3\tau_1 \wedge \phi + *\tau_3, \\ d*\phi &= 4\tau_1 \wedge *\phi + \tau_2 \wedge \phi, \end{aligned}$$

for suitable forms  $\tau_p \in \Omega^p$ . In terms of the  $G_2$  invariant splittings of the exterior algebra,  $\tau_0 \in \Omega_1^0$ ,  $\tau_1 \in \Omega_7^1$ ,  $\tau_2 \in \Omega_{14}^2$ ,  $\tau_3 \in \Omega_{27}^3$ . The notation  $\Omega_d^p$  indicates the space of  $p$ -forms taking values in the  $d$ -dimensional  $G_2$  irreducible subspace  $\Lambda_d^p \subset \Lambda^p$ . The one-form  $\tau_1$  is also known as the *Lee form* of the  $G_2$  manifold. The forms  $\tau_0, \tau_1, \tau_2, \tau_3$  correspond to the Fernández-Gray classes [10] as follows

$$\tau_0 \leftrightarrow \mathcal{X}_1, \quad \tau_2 \leftrightarrow \mathcal{X}_2, \quad \tau_3 \leftrightarrow \mathcal{X}_3, \quad \tau_1 \leftrightarrow \mathcal{X}_4.$$

When we speak of the intrinsic torsion  $\tau$  of a  $G_2$  structure we mean the form of mixed degree  $\tau = \tau_0 + \tau_1 + \tau_2 + \tau_3$  fixed by the fundamental three-form as above.

In particular,  $G_2$ -manifolds in the class  $\mathfrak{X}_1$  are characterized by the conditions  $\tau_1 = \tau_2 = \tau_3 = 0$  and are called nearly parallel  $G_2$ -manifolds. It is well known that these spaces are Einstein with positive scalar curvature. From this it follows that  $\tau_0$  is constant [13, 14].

A  $G_2$ -manifold in the Fernández-Gray class  $\mathfrak{X}_1 + \mathfrak{X}_4$  satisfies  $\tau_2 = \tau_3 = 0$ . The structure equations for this case reduce to

$$\begin{aligned} d\phi &= \tau_0 * \phi + 3\tau_1 \wedge \phi, \\ d*\phi &= 4\tau_1 \wedge * \phi, \end{aligned}$$

from which one infers

$$(3.2) \quad \begin{aligned} d^2\phi &= (d\tau_0 + \tau_0\tau_1) \wedge * \phi + 3d\tau_1 \wedge \phi, \\ d^2*\phi &= 4d\tau_1 \wedge * \phi. \end{aligned}$$

The latter equation implies that the component  $(d\tau_1)_7 \in \Omega_7^2$  vanishes. Using this in the first equation, one deduces that the complementary component  $(d\tau_1)_{14} \in \Omega_{14}^2$  also vanishes. Thus we recover the fact (see [23]) that  $G_2$  structures in this class are, locally, conformal to a nearly parallel structure. Observe that equations (3.2) furthermore give us  $d\tau_0 + \tau_0\tau_1 = 0$ . Now our Lemma 3 applies with  $\phi = \tau_0$  and  $\alpha = \tau_1$ . The connectedness of  $M$  leads to the conclusion in the form of this

**Theorem 4.** *Suppose  $M$  is a 7-dimensional manifold with a  $G_2$ -structure  $\phi$  in the Fernández-Gray class  $\mathfrak{X}_1 + \mathfrak{X}_4$ . Then  $\phi$  is either of class  $\mathfrak{X}_4$ , in which case  $(M, \phi)$  is locally conformal to a parallel  $G_2$ -manifold, or  $(M, \phi)$  is conformal to a nearly parallel  $G_2$ -manifold.*

#### 4. THE 6-DIMENSIONAL ALMOST HERMITIAN CASE

An almost Hermitian manifold is a Riemannian manifold  $(M^{2m}, g)$  equipped with an orthogonal almost complex structure  $J$ . The metric and the almost complex structure then define the fundamental two-form of the almost Hermitian structure:

$$\omega(X, Y) := g(JX, Y).$$

As opposed to the  $G_2$  case above and the case of  $SU(3)$  below, the components of the intrinsic torsion of a Hermitian structure cannot all be identified with differential forms [27]. Instead, the intrinsic torsion is detected by  $d\omega$  along with the Nijenhuis tensor  $N_J$ .

From now on  $m$  is taken to be at least 3.

**4.1. The Nijenhuis Tensor.** For an almost complex structure  $J$  the Nijenhuis tensor measures the failure of the eigenspaces  $J$  in the complexified tangent space to be involutive. Let  $\pi'(X) = \frac{1}{2}(X - iJX)$  and  $\pi'' = \frac{1}{2}(X + iJX)$  be the projections to the  $i$ -eigenspace  $T'$  and the  $-i$ -eigenspace  $T''$ , respectively. We set

$$\begin{aligned} N_J(X, Y) &:= \pi'[\pi''X, \pi''Y] + \pi''[\pi'X, \pi'Y] \\ &= \frac{1}{4}([X, Y] - [JX, JY] + J[JX, Y] + J[X, JY]). \end{aligned}$$

Using the metric we obtain an algebraically equivalent 3-tensor

$$N_J(X, Y; Z) = g(N_J(X, Y), Z),$$

with the property  $N_J(JX, Y; Z) = N_J(X, JY; Z) = N_J(X, Y; JZ)$ . Equivalently,

$$N_J \in \llbracket \Lambda^{(2,0)} \otimes \Lambda^{(1,0)} \rrbracket.$$

See for instance [9] for an explanation of the notation.

The space  $\llbracket \Lambda^{(3,0)} \rrbracket$  is a subspace of  $\llbracket \Lambda^{(2,0)} \otimes \Lambda^{(1,0)} \rrbracket$  in the natural way. The projection  $\llbracket \Lambda^{(2,0)} \otimes \Lambda^{(1,0)} \rrbracket \rightarrow \llbracket \Lambda^{(3,0)} \rrbracket$  is given simply by skew-symmetrization. Write  $V$  for the orthogonal complement of  $\llbracket \Lambda^{(3,0)} \rrbracket$  in  $\llbracket \Lambda^{(2,0)} \otimes \Lambda^{(1,0)} \rrbracket$ . Then we may split the Nijenhuis tensor accordingly

$$N_J = N_J^{3,0} + N_J^V.$$

One may now deduce that

$$(4.1) \quad (d\omega)^{3,0}(X, Y, Z) = 3g(N_J^{3,0}(X, Y), JZ) = 3N_J^{3,0}(X, Y, JZ).$$

The structure equations for an almost Hermitian manifold now can be written

$$(4.2) \quad d\omega = -3J_{(1)}N_J^{3,0} + 2\sigma_1 \wedge \omega + \sigma_3, \quad N_J = N_J^{3,0} + N_J^V.$$

The first equation here employs the conventions  $(J_{(1)}\alpha)(X, Y, \dots) := -\alpha(JX, Y, \dots)$ , see [24]. This action of the complex structure  $J$  on differential forms is, generally speaking, distinct from the usual action given by  $(J\alpha)(X_1, \dots, X_p) := (-1)^p \alpha(JX_1, \dots, JX_p)$ .

The Gray-Hervella classes of an almost Hermitian manifold are in the following correspondence with the components in (4.2)

$$N_J^{3,0} \leftrightarrow \mathcal{W}_1, \quad N_J^V \leftrightarrow \mathcal{W}_2, \quad \sigma_3 \leftrightarrow \mathcal{W}_3, \quad \sigma_1 \leftrightarrow \mathcal{W}_4.$$

Almost Hermitian manifolds in the class  $\mathcal{W}_1$ , called nearly Kähler manifolds, are characterized by the conditions  $N_J^V = \sigma_1 = \sigma_3 = 0$  or equivalently, by demanding that the covariant derivative of the almost complex structure with respect to the Levi-Civita connection be skew-symmetric,  $(\nabla_X^g J)X = 0$  [15].

*Remark 5.* Almost Hermitian manifolds in the class  $\mathcal{W}_1 + \mathcal{W}_3 + \mathcal{W}_4$  are characterized by  $N_J^V = 0$ , i.e. the Nijenhuis tensor is totally skew-symmetric. This amounts to the existence of a linear connection preserving the almost Hermitian structure with totally skew-symmetric torsion [12]. In dimension 6, this class can also be characterized by the property that the Nijenhuis tensor is either everywhere non-degenerate (of constant signature) or vanishes identically [5]. These manifolds are called quasi-integrable and investigated in details in [5].

For an arbitrary one-form the relation

$$(d\theta)(X, Y) - (d\theta)(JX, JY) - d(J\theta)(JX, Y) + Jd(J\theta)(JX, Y) = 4g(N_J(X, Y), J(J\theta)^\#)$$

holds. Writing  $d\theta^{2,0} = \frac{1}{2}(d\theta - Jd\theta)$  for the projection of  $d\theta$  to  $\llbracket \Lambda^{(2,0)} \rrbracket$  we have

**Lemma 6.** *Suppose  $(g, J)$  is an almost Hermitian structure in class  $\mathcal{W}_1 + \mathcal{W}_3 + \mathcal{W}_4$ . Let  $\theta$  be a one-form and write  $\theta' := J\theta$ . Then*

$$(d\theta)^{2,0} + J_{(1)}(d\theta')^{2,0} = \frac{2}{3}\theta' \lrcorner (d\omega)^{3,0}$$

**4.2.  $SU(3)$ -structures.** A 6-dimensional manifold with an  $SU(3)$ -structure comes equipped with data  $(g, J, \omega, \psi_+, \psi_-)$  invariant with respect to the action of  $SU(3)$ . Here  $g$  is a Riemannian metric,  $J$  is an almost complex structure,  $\omega$  the fundamental two-form and  $\psi_+$  and  $\psi_-$  are three-forms such that  $\Psi := \psi_+ + i\psi_-$  is a complex  $(3,0)$ -form. These invariant tensors are not independent, in fact the triple  $(\omega, \psi_+, \psi_-)$  with  $\psi_+ + i\psi_-$  decomposable and compatible with  $\omega$  by means of the equations below, defines both  $g$  and  $J$ , see [19]. Clearly the triple  $(g, \psi_+, \psi_-)$  will do the same. We choose a normalization with the following relations

$$(4.3) \quad \begin{aligned} \omega(X, JY) &= g(X, Y), \\ \omega \wedge \psi_+ &= 0 = \omega \wedge \psi_-, \\ 3\psi_+ \wedge \psi_- &= 2\omega^3 = 12 \operatorname{vol}_g, \\ *\omega &= \frac{1}{2}\omega^2, \quad *\psi_+ = \psi_-, \quad J\psi_+ = -\psi_-. \end{aligned}$$

**4.2.1. Torsion classes and structure equations.** Under the action of  $SU(3)$ ,  $\Lambda^{(3,0)} = \mathbb{C}$  and  $V \otimes \mathbb{C} \cong \Lambda_0^{(1,1)}$ . This means that  $V \cong 2\mathfrak{su}(3)$  and  $\llbracket \Lambda^{(3,0)} \rrbracket \cong 2\mathbb{R}$ . Moreover, for an  $SU(3)$ -structure  $(\omega, \psi_\pm)$ , the components of the Nijenhuis tensor can be computed from components of  $(d\omega, d\psi_\pm)$ . In fact there are algebraic correspondences

$$\begin{aligned} N_J^{(3,0)} &\leftrightarrow (d\omega)^{(3,0)+(0,3)} \leftrightarrow \left( (d\psi_+)^{(0,0)}, (d\psi_-)^{(0,0)} \right), \\ N_J^V &\leftrightarrow \left( (d\psi_+)_0^{(2,2)}, (d\psi_-)_0^{(2,2)} \right). \end{aligned}$$

The first arrow is given by equation (4.1), the second is described below. The remaining one has a similar description which we will not need here.

The  $SU(3)$ -structure function  $\nabla^g \Psi$  is completely determined by the exterior derivatives of the three forms  $\omega, \psi_+$  and  $\psi_-$ . These may be written as

$$(4.4) \quad \begin{aligned} d\omega &= 3(\sigma_0^+ \psi_+ - \sigma_0^- \psi_-) + 2\sigma_1^+ \wedge \omega + \sigma_3, \\ d\psi_+ &= -2\sigma_0^- \omega^2 + 3\sigma_1^+ \wedge \psi_+ - \sigma_1^- \wedge \psi_- + \sigma_2^+ \wedge \omega, \\ d\psi_- &= -2\sigma_0^+ \omega^2 + 3\sigma_1^+ \wedge \psi_- + \sigma_1^- \wedge \psi_+ + \sigma_2^- \wedge \omega. \end{aligned}$$

Here  $\sigma_p^\pm$  are  $p$ -forms and  $\sigma_3$  is a three-form. They correspond roughly to the classes  $\mathcal{W}_1^+, \mathcal{W}_1^-, \mathcal{W}_4, \mathcal{W}_5, \mathcal{W}_2^+, \mathcal{W}_2^-$ , and  $\mathcal{W}_3$  of [8], respectively (see also [4]). These determine the Gray-Hervella classes of the underlying almost Hermitian structure in the obvious way.

*Remark 7.* The one-form  $\sigma_1^+$  is, in fact, the Lee form of the almost Hermitian structure. In contrast  $\sigma_1^-$  is conformally invariant. Therefore  $\sigma_1^-$  does not really correspond to the class  $\mathcal{W}_5$  but rather to “ $3\mathcal{W}_4 + 2\mathcal{W}_5$ ”. This choice for the one-forms was introduced by Martín Cabrera [24].

4.2.2. *A transformation.* Set  $\lambda := \sigma_0^+ + i\sigma_0^-$  and  $\Lambda := |\lambda|$ . In neighbourhoods with  $\lambda$  non-vanishing an argument  $\varphi := \arg(\lambda) := \arctan\left(\frac{\sigma_0^-}{\sigma_0^+}\right)$  may be chosen. We then set

$$\begin{aligned} \tilde{\omega} &:= \Lambda^2 \omega, \\ \tilde{\psi}_+ &:= \Lambda^2 (\sigma_0^+ \psi_+ - \sigma_0^- \psi_-), \\ \tilde{\psi}_- &:= \Lambda^2 (\sigma_0^- \psi_+ + \sigma_0^+ \psi_-). \end{aligned}$$

This gives the somewhat simpler structure equations

$$\begin{aligned} d\tilde{\omega} &:= 3\tilde{\psi}_+ + 2\tilde{\sigma}_1^+ \wedge \omega + \tilde{\sigma}_3, \\ d\tilde{\psi}_+ &:= 3\tilde{\sigma}_1^+ \wedge \tilde{\psi}_+ - \tilde{\sigma}_1^- \wedge \tilde{\psi}_- + \tilde{\sigma}_2^+ \wedge \tilde{\omega}, \\ d\tilde{\psi}_- &:= -2\tilde{\omega}^2 + 3\tilde{\sigma}_1^+ \wedge \tilde{\psi}_- + \tilde{\sigma}_1^- \wedge \tilde{\psi}_+ + \tilde{\sigma}_2^- \wedge \tilde{\omega}. \end{aligned}$$

where

$$\begin{aligned} \tilde{\sigma}_1^+ &:= \sigma_1^+ + \Lambda^{-1} d\Lambda, & \tilde{\sigma}_1^- &:= \sigma_1^- - d\varphi, \\ \tilde{\sigma}_2^+ &:= \sigma_0^+ \sigma_2^+ - \sigma_0^- \sigma_2^-, & \tilde{\sigma}_2^- &:= \sigma_0^- \sigma_2^+ + \sigma_0^+ \sigma_2^-, \\ \tilde{\sigma}_3 &:= \Lambda^2 \sigma_3. \end{aligned}$$

In particular, the structure equations of a nearly Kähler 6-manifold can always be put on the form [18, 28]

$$d\omega = 3\psi_+, \quad d\psi_- = -2\omega^2.$$

It is well known that these spaces are Einstein with positive scalar curvature [16].

## 5. LOCALLY CONFORMALLY NEARLY KÄHLER 6-FOLDS

For a 6-dimensional almost Hermitian manifold in the class  $\mathcal{W}_1 + \mathcal{W}_4$  a (possibly local) choice of trivialization  $(\psi_+, \psi_-)$  allows us to write the structure equations (4.2), (4.4) as

$$(5.1) \quad \begin{aligned} d\omega &= 3(\sigma_0^+ \psi_+ - \sigma_0^- \psi_-) + 2\sigma_1^+ \wedge \omega, \\ d\psi_+ &= -2\sigma_0^- \omega^2 + 3\sigma_1^+ \wedge \psi_+ - \sigma_1^- \wedge \psi_-, \\ d\psi_- &= -2\sigma_0^+ \omega^2 + 3\sigma_1^+ \wedge \psi_- + \sigma_1^- \wedge \psi_+. \end{aligned}$$

Differentiating each equation yields

$$(5.2) \quad 0 = 3(d\sigma_0^+ + \sigma_0^+ \sigma_1^+ - \sigma_0^- \sigma_1^-) \psi_+ - 3(d\sigma_0^- + \sigma_0^- \sigma_1^+ + \sigma_0^+ \sigma_1^-) \psi_- + 2d\sigma_1^+ \omega,$$

$$(5.3) \quad 0 = -2(d\sigma_0^- + \sigma_0^- \sigma_1^+ + \sigma_0^+ \sigma_1^-) \omega^2 + 3d\sigma_1^+ \psi_+ - d\sigma_1^- \psi_-,$$

$$(5.4) \quad 0 = -2(d\sigma_0^+ + \sigma_0^+ \sigma_1^+ - \sigma_0^- \sigma_1^-) \omega^2 + 3d\sigma_1^+ \psi_- + d\sigma_1^- \psi_+.$$

These have the following immediate consequences. Equation (5.2) shows that  $d\sigma_1^+$  is a  $(2, 0) + (0, 2)$  form as a linear combination of one-forms contracted with  $\psi_+$  and  $\psi_-$ . Using standard identities such as  $J(\sigma \wedge \psi_+) =$

$\sigma \wedge \psi_-$  for an arbitrary two-form  $\sigma$  and  $J(\sigma \wedge \omega) = \sigma \wedge \omega$  for a  $(1,1)$ -form, as well as  $*(\theta \wedge \psi_-) = \theta \lrcorner \psi_+ = (J\theta) \lrcorner \psi_-$  for a one-form  $\theta$ , leads to the equivalent set of equations:

$$(5.5) \quad d\sigma_0^+ + \sigma_0^+ \sigma_1^+ - \sigma_0^- \sigma_1^- = J(d\sigma_0^- + \sigma_0^- \sigma_1^+ + \sigma_0^+ \sigma_1^-),$$

$$(5.6) \quad d\sigma_1^+ = 3(d\sigma_0^- + \sigma_0^- \sigma_1^+ + \sigma_0^+ \sigma_1^-) \lrcorner \psi_+,$$

$$(5.7) \quad (d\sigma_1^-)^{2,0} = 7(d\sigma_0^- + \sigma_0^- \sigma_1^+ + \sigma_0^+ \sigma_1^-) \lrcorner \psi_-.$$

**Lemma 8.** *Suppose  $(M^6, \omega, J)$  is an almost Hermitian manifold in the class  $\mathcal{W}_1 + \mathcal{W}_4$ . Then the Lee form is closed if and only if  $(\omega, J)$  is either globally conformal to a nearly Kähler structure  $(\omega', J')$  on  $M$  or locally conformally equivalent to a Kähler structure.*

*Proof.* Suppose the Lee-form is closed  $d\sigma_1^+ = 0$ . Locally, we pick a smooth trivialisation  $(\psi_+, \psi_-)$  of  $[\Lambda^{(3,0)}]$ . Then, locally, equations (5.6) and (5.5) show that

$$\begin{aligned} d\sigma_0^+ + \sigma_0^+ \sigma_1^+ - \sigma_0^- \sigma_1^- &= 0, \\ d\sigma_0^- + \sigma_0^- \sigma_1^+ + \sigma_0^+ \sigma_1^- &= 0, \end{aligned}$$

whence

$$d((\sigma_0^+)^2 + (\sigma_0^-)^2) + ((\sigma_0^+)^2 + (\sigma_0^-)^2)(2\sigma_1^+) = 0.$$

However,

$$\phi := (\sigma_0^+)^2 + (\sigma_0^-)^2 = \frac{1}{9} \|d\omega^{3,0}\|^2$$

is a globally well-defined, smooth function, and  $\alpha := 2\sigma_1^+$  is closed. So Lemma 3 applies and we conclude that  $\|d\omega^{3,0}\|$  is either non-zero everywhere, or it vanishes at all points.  $\square$

*Remark 9.* Note that in dimension 6 the Nijenhuis tensor is either everywhere non-degenerate (of constant signature) or vanishes identically not only for  $\mathcal{W}_1 + \mathcal{W}_4$ , but for the whole class  $\mathcal{W}_1 + \mathcal{W}_3 + \mathcal{W}_4$  [5].

**Theorem 10.** *Let  $M$  be a 6-dimensional manifold with an almost Hermitian structure  $(\omega, J)$  in the Gray-Hervella class  $\mathcal{W}_1 + \mathcal{W}_4$ . Then either  $(\omega, J)$  is locally conformally equivalent to Kähler structure on  $M$ , or  $(\omega, J)$  is a conformal transformation of a nearly Kähler structure.*

*Proof.* Write  $M$  as a disjoint union  $M_0 \cup M_1$  where

$$M_0 := \{x \in M : (d\omega)^{3,0} = 0\}, \quad M_1 := \{x \in M : (d\omega)^{3,0} \neq 0\}.$$

On the open submanifold  $M_1$  there is a canonical choice of trivialization of  $[\Lambda^{(3,0)}]$  given by taking  $\psi_+ = (d\omega)^{3,0}$ . After a suitable transformation (as in section 4.2.2) we obtain the structure equations

$$\begin{aligned} (5.8) \quad d\tilde{\omega} &= 3\tilde{\psi}_+ + 2\tilde{\sigma}_1^+ \wedge \tilde{\omega}, \\ d\tilde{\psi}_+ &= 3\tilde{\sigma}_1^+ \wedge \tilde{\psi}_+ - \tilde{\sigma}_1^- \wedge \tilde{\psi}_-, \\ d\tilde{\psi}_- &= -2\tilde{\omega}^2 + 3\tilde{\sigma}_1^+ \wedge \tilde{\psi}_- + \tilde{\sigma}_1^- \wedge \tilde{\psi}_+. \end{aligned}$$

Equations (5.5), (5.6) and (5.7) then become

$$\begin{aligned} (5.9) \quad \tilde{\sigma}_1^+ &= J\tilde{\sigma}_1^-, \\ d\tilde{\sigma}_1^+ &= 3\tilde{\sigma}_1^- \lrcorner \tilde{\psi}_+, \\ (d\tilde{\sigma}_1^-)^{2,0} &= 7\tilde{\sigma}_1^- \lrcorner \tilde{\psi}_-. \end{aligned}$$

Using Lemma 6 with  $\theta = \tilde{\sigma}_1^+$ ,  $\theta' = \tilde{\sigma}_1^-$ , and the identity  $J_{(1)}(\sigma \lrcorner \psi_{\pm}) = (J\sigma) \lrcorner \psi_{\pm} = \mp \sigma \lrcorner \psi_{\mp}$  valid for all one-forms  $\sigma$ , we get

$$d\tilde{\sigma}_1^+ - J_{(1)}(d\tilde{\sigma}_1^-)^{2,0} = -2\tilde{\sigma}_1^+ \lrcorner \tilde{\psi}_- = -2\tilde{\sigma}_1^- \lrcorner \tilde{\psi}_+.$$

This is only compatible with the relations (5.9) if  $\tilde{\sigma}_1^- \lrcorner \tilde{\psi}_+ = 0$ . Therefore  $\tilde{\sigma}_1^+ = \tilde{\sigma}_1^- = 0$  and the original one-forms  $\sigma_1^{\pm}$  are, in fact, exact on  $M_1$ . Moreover, on the interior  $M_0^o$  of  $M_0$ ,  $d\omega = 2\sigma_1^+ \wedge \omega$ , so  $d\sigma_1^+|_{M_0^o} = 0$  also holds.

So the set of points at which  $d\sigma_1^+ \neq 0$ , which clearly is open, is the common boundary of two open sets in  $M$ , at least one of which is non-empty. Therefore  $d\sigma_1^+ = 0$  on all of  $M$  and Lemma 8 completes the proof.  $\square$

## 6. PROOF OF THEOREM 1 AND THEOREM 2

Theorem 4 and Theorem 10 show that the Riemannian manifold  $(M, g)$  is globally conformal to an Einstein space of positive scalar curvature. Further, if the scalar curvature is constant then the Obata Theorem (see the proof in [22]) tells us that the conformal transformation making the metric Einstein is trivial, or else  $(M, g)$  is the standard sphere.  $\square$

## 7. CURVATURE CLASSIFICATION

The Riemannian curvature tensor of a nearly Kähler 6-manifolds or a nearly parallel  $G_2$ -manifold is especially simple. In fact, viewing curvature tensors as bundle endomorphisms  $R: \Lambda^2 \rightarrow \Lambda^2$ , the curvature splits as

$$(7.1) \quad R^g = R^{\mathfrak{g}} + \frac{s_g}{2n(n-1)} \text{Id}_{\Lambda^2}.$$

where  $R^{\mathfrak{g}}$ , formally, is the curvature tensor of a space with holonomy algebra  $\mathfrak{g}$  and  $n$  is the dimension of the underlying space. This formula is reminiscent of the curvature formula for a Riemannian manifold with holonomy  $Sp(n)Sp(1)$  of [27]. In the cases of concern,  $\mathfrak{g}$  and  $n$  are equal to  $\mathfrak{su}(3)$  and 6 for nearly Kähler and  $\mathfrak{g}_2$  and 7 for nearly parallel  $G_2$ . In either situation the tensor  $R^{\mathfrak{g}}$  takes values in a  $G$ -irreducible subspace of the space of algebraic Weyl tensors, i.e., algebraic curvature tensors with vanishing Ricci contraction. Standard identities [3] now make it possible to deduce the form of the curvature tensor for an almost Hermitian or  $G_2$  space of type  $\mathcal{W}_1 + \mathcal{W}_4$ , or  $\mathcal{X}_1 + \mathcal{X}_4$ , respectively. The conformal invariance of the Weyl curvature implies that a manifold in the more general classes are Einstein if and only if their curvature is of the same form of nearly parallel structures.

**Theorem 11.** (a) *Suppose  $(M, \phi)$  is a  $G_2$  manifold of strict type  $\mathcal{X}_1 + \mathcal{X}_4$  such that the associated metric  $g$  is complete and Einstein. Then  $(M, g)$  is isometric to either the sphere, the hyperbolic space or the euclidean space equipped with a constant curvature metric. Furthermore, the  $G_2$  structure  $\phi$  is a conformal change of a suitable restriction of the standard nearly parallel structure on the 7 dimensional sphere.*  
(b) *Suppose  $(M, \omega, J)$  is an almost Hermitian 6-manifold of strict type  $\mathcal{W}_1 + \mathcal{W}_4$  such that the associated metric  $g$  is complete and Einstein. Then  $(M, g)$  is isometric to either the sphere, hyperbolic space or euclidean space equipped with a constant curvature metric. Furthermore, the almost Hermitian structure  $(\omega, J)$  is a conformal change of a suitable restriction of the standard nearly Kähler structure on the 6 dimensional sphere.*

*Proof.* This is an immediate consequence of Theorem 4, Theorem 10, uniqueness of the nearly parallel structure compatible with the round metric on the 7-sphere, uniqueness of the nearly Kähler structure compatible with the round metric on the 6-sphere, see Friedrich [11] and the Main Theorem of [21].  $\square$

**Corollary 12.** *Suppose  $M$  is a  $G_2$  manifold or almost Hermitian 6-manifold of strict type  $\mathcal{X}_1 + \mathcal{X}_4$  or  $\mathcal{W}_1 + \mathcal{W}_4$ , respectively. Assume that the Riemannian curvature of  $M$  is of the form (7.1). Then  $M$  has constant sectional curvature and in particular  $R^{\mathfrak{g}} = 0$ .*

## REFERENCES

- [1] Ilka Agricola and Thomas Friedrich. Geometric structures of vectorial type. 1
- [2] Ilka Agricola and Thomas Friedrich. On the holonomy of connections with skew-symmetric torsion. *Math. Ann.*, 328(4):711–748, 2004. 1
- [3] Arthur L. Besse. *Einstein manifolds*, volume 10 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1987. 7
- [4] Gil Bor and Luis Hernández Lamóneda. Bochner formulae for orthogonal  $G$ -structures on compact manifolds. *Differential Geom. Appl.*, 15(3):265–286, 2001. 5
- [5] Robert L. Bryant. Remarks on the geometry of almost complex 6-manifolds. 4, 6
- [6] Robert L. Bryant. Some remarks on  $G_2$ -structures. 1, 2
- [7] Jean-Baptiste Butruille. Espace de twisteurs réduit d’une variété presque hermitienne de dimension 6. 1

- [8] Simon Chiossi and Simon Salamon. The intrinsic torsion of  $SU(3)$  and  $G_2$  structures. In *Differential geometry, Valencia, 2001*, pages 115–133. World Sci. Publishing, River Edge, NJ, 2002. 1, 5
- [9] Maria Falcitelli, Angela Farinola, and Simon Salamon. Almost-Hermitian geometry. *Differential Geom. Appl.*, 4(3):259–282, 1994. 1, 3
- [10] Marisa Fernández and Alfred Gray. Riemannian manifolds with structure group  $G_2$ . *Ann. Mat. Pura Appl. (4)*, 132:19–45 (1983), 1982. 1, 2
- [11] Thomas Friedrich. Nearly Kaehler and nearly parallel  $G_2$ -structures on spheres. 7
- [12] Thomas Friedrich and Stefan Ivanov. Parallel spinors and connections with skew-symmetric torsion in string theory. *Asian J. Math.*, 6(2):303–335, 2002. 1, 4
- [13] Thomas Friedrich, Ines Kath, Andrei Moroianu, and Uwe Semmelmann. On nearly parallel  $G_2$ -structures. *J. Geom. Phys.*, 23(3-4):259–286, 1997. 3
- [14] Krzysztof Galicki and Simon Salamon. Betti numbers of 3-Sasakian manifolds. *Geom. Dedicata*, 63(1):45–68, 1996. 3
- [15] Alfred Gray. Nearly Kähler manifolds. *J. Differential Geometry*, 4:283–309, 1970. 4
- [16] Alfred Gray. Weak holonomy groups. *Math. Z.*, 123:290–300, 1971. 5
- [17] Alfred Gray and Luis M. Hervella. The sixteen classes of almost Hermitian manifolds and their linear invariants. *Ann. Mat. Pura Appl. (4)*, 123:35–58, 1980. 1
- [18] Nigel Hitchin. Stable forms and special metrics. In *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, volume 288 of *Contemp. Math.*, pages 70–89. Amer. Math. Soc., Providence, RI, 2001. 5
- [19] Nigel J. Hitchin. The moduli space of special Lagrangian submanifolds. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 25(3-4):503–515 (1998), 1997. Dedicated to Ennio De Giorgi. 4
- [20] Stefan Ivanov, Maurizio Parton, and Paolo Piccinni. Locally conformal parallel  $g_2$  and  $spin7$  manifolds. *Math. Res. Lett.*, 13(2):167–177, 2006. 1
- [21] Wolfgang Kühnel. Conformal transformations between Einstein spaces. In *Conformal geometry (Bonn, 1985/1986)*, Aspects Math., E12, pages 105–146. Vieweg, Braunschweig, 1988. 7
- [22] John M. Lee and Thomas H. Parker. The Yamabe problem. *Bull. Amer. Math. Soc. (N.S.)*, 17(1):37–91, 1987. 7
- [23] Francisco Martín Cabrera. On Riemannian manifolds with  $G_2$ -structure. *Boll. Un. Mat. Ital. A (7)*, 10(1):99–112, 1996. 1, 3
- [24] Francisco Martín Cabrera. Special almost Hermitian geometry. *J. Geom. Phys.*, 55(4):450–470, 2005. 1, 4, 5
- [25] Francisco Martín Cabrera, Maria D. Monar, and Andrew F. Swann. Classification of  $G_2$ -structures. *J. London Math. Soc. (2)*, 53(2):407–416, 1996. 1
- [26] Francisco Martín Cabrera and Andrew Swann. Curvature of (special) almost Hermitian manifolds. 2
- [27] Simon Salamon. *Riemannian geometry and holonomy groups*, volume 201 of *Pitman Research Notes in Mathematics Series*. Longman Scientific & Technical, Harlow, 1989. 3, 7
- [28] Simon Salamon. Almost parallel structures. In *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, volume 288 of *Contemp. Math.*, pages 162–181. Amer. Math. Soc., Providence, RI, 2001. 5
- [29] Misha Verbitsky. An intrinsic volume functional on almost complex 6-manifolds and nearly Kaehler geometry. 1

(Cleyton) HUMBOLDT-UNIVERSITÄT ZU BERLIN, INSTITUT FÜR MATHEMATIK, UNTER DEN LINDEN 6, D-10099 BERLIN, GERMANY

*E-mail address:* cleyton@mathematik.hu-berlin.de

(Ivanov) UNIVERSITY OF SOFIA "ST. KL. OHRIDSKI", FACULTY OF MATHEMATICS AND INFORMATICS,, BLVD. JAMES BOURCHIER 5,, 1164 SOFIA, BULGARIA

*E-mail address:* ivanovsp@fmi.uni-sofia.bg